

ETS-Related Gene (ERG) and Friend leukemia integration – 1 (FLI-1) Transcription Factors in the Precision Treatment of Pulmonary Arterial Hypertension and Pulmonary Fibrosis

Titilope Olanipekun MD¹, Nicolas Bakinde MD¹, Kelsey Paul-Stubbs MD², Ajose Taiwo MD¹, Sharif Morsalin PhD², Samir Yezdani MD², Veena N Rao PhD², and E. Shyam P Reddy PhD^{2*}

¹Internal Medicine Program, Morehouse School of Medicine, Atlanta, GA 30303

²Cancer Biology Program, Department of OB/GYN, Morehouse School of Medicine, Georgia Cancer Center for Excellence, Grady Memorial Hospital, Atlanta, GA 30303

*Corresponding author: E. Shyam P. Reddy, PhD, Professor and Director, Cancer Biology Program, Department of OB/GYN, Morehouse School of Medicine, Georgia Cancer Center for Excellence, RM 10C009, Grady Memorial Hospital, 80 Jesse Hill Jr. Drive, Atlanta, GA 30303, USA; Tel: 404-489-9991; E mail: ereddy@msm.edu

Article Type: Review, **Submission Date:** 31 May 2019, **Accepted Date:** 10 June 2019, **Published Date:** 26 June 2019.

Citation: Titilope Olanipekun MD, Nicolas Bakinde MD, Kelsey Paul-Stubbs MD, Ajose Taiwo MD, Sharif Morsalin PhD, et al. (2019) ETS-Related Gene (ERG) and Friend leukemia integration – 1 (FLI-1) Transcription Factors in the Precision Treatment of Pulmonary Arterial Hypertension and Pulmonary Fibrosis. *J Can Epi Treat* 3(1): 1-4. doi: <https://doi.org/10.24218/jcet.2019.21>.

Copyright: © 2019 Titilope Olanipekun MD, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Pulmonary arterial hypertension (PAH) is a chronic debilitating cardiopulmonary disease characterized by abnormal remodeling of peripheral lung vasculature resulting in progressive vasoconstriction. Idiopathic pulmonary fibrosis (IPF) is a chronic, progressive, and irreversible disease that is often associated with significant morbidity and poor quality of life. The prognosis of PAH and IPF is poor and currently available medications focus on relieving symptoms and slowing down progression. Hence, there is a clear necessity to develop new therapies. ETS-related genes and Friend leukemia integration-1 (FLI-1) are transcription factors involved in angiogenesis, cellular homeostasis, vascular remodeling, and the genetic regulation of inflammation, apoptosis, and fibrosis seen in PAH and IPF. Simultaneous small-interfering-RNA (siRNA) knockdown of ERG and FLI-1 in human pulmonary artery endothelial cells (HPAEC) and human pulmonary microvasculature endothelial cells (HPMEC) has been associated with up-regulation of pro-inflammatory genes and interferon (IFN) pathway-related genes. Notably, the endothelium in normal lungs has also been shown to have high levels of nuclear ERG and Fli-1 compared to significantly lower levels in diseased lungs. Recently, ERG upregulation was found to promote liver homeostasis by regulating canonical TGFβ1-SMAD signaling and promoting the SMAD1 pathway while repressing SMAD3 activity. Improvement in pulmonary fibrosis through medications that suppress the TGF-β1/Smad3 pathway has also been a subject of study. In this review, we hypothesize that targeting the regulation of ERG, FLI-1 and ERG-mediation

of TGF-β1/Smad3 signaling may be a promising therapeutic strategy in PAH and IPF.

Keyword: Pulmonary Diseases, ETS genes, Signaling pathways, Targeted therapy, small molecules, Smad3 inhibitors, ERG/Fli-1 inducers.

Introduction

Pulmonary arterial hypertension (PAH) is a chronic debilitating cardiopulmonary disease characterized by abnormal remodeling of peripheral lung vasculature resulting in progressive vasoconstriction [1]. PAH is associated with an elevated pulmonary arterial pressure (> 25 mm Hg at rest to > 30 mm Hg with exercise) and if left untreated leads to right-heart hypertrophy, and consequently heart failure [2]. PAH could be classified as idiopathic (formerly known as primary pulmonary hypertension) or associated with autoimmune, infectious, vascular, metabolic, and congenital disorders [1]. The underlying pathology of PAH involves the recruitment of inflammatory cells with release of cytokines which enhance cell proliferation and elastin fibers degradation [3]. Histologically, PAH is characterized by intimal fibrosis, increased medial thickness, predominance of complex plexiform lesions resulting in vasoconstriction and pulmonary arteriolar occlusion [4].

Idiopathic pulmonary fibrosis (IPF) is a chronic, progressive, and irreversible disease that is often associated with significant morbidity and poor quality of life [5]. Till date, the cause of IPF is unknown and the pathogenesis is not completely clear [6]. However, studies have shown that the fibrosis is driven by the

abnormal activation of alveolar epithelial cells (AECs) by TGF- β 1, which enhance the formation of fibroblasts and myofibroblasts, through the proliferation of resident mesenchymal cells and stimulation of the epithelial to mesenchymal transition (EMT) [6].

The disease mechanism in PAH and IPF has been the target of numerous therapeutic interventions [7]. The current mainstay of PAH treatment essentially involves promoting vasodilation, inhibiting prothrombotic and inflammatory processes. Similarly, in IPF, treatment is mainly geared towards relieving symptoms as much as possible and slowing down the progression. The prognosis of PAH and IPF is poor [1,6], hence, there is a clear necessity to develop new therapies.

Recent evidence suggests that there are families of transcription factors (ETS-related gene (ERG) and Friend leukemia integration-1 (FLI1) involved in the angiogenesis, cellular homeostasis, vascular remodeling, and the genetic regulation of inflammation, apoptosis, and fibrosis seen in PAH and IPF [8-10]. Improved understanding of the genetic modulation of these processes has revealed novel therapeutic strategies. In this review, we will discuss the role of the transcription factors in the pathogenesis of PAH and IPF, and they can hypothetically contribute to precision and targeted therapy.

FLI1 and ERG in the Modulation of Vascular Homeostasis, Inflammation and Tissue Fibrosis

ERG and human FLI1 both belong to the ETS family of transcriptional factors and were discovered by Drs. Reddy and Rao [11,12]. ETS transcription factors share an 85 amino acid DNA binding domain (ETS domain) that binds to a DNA core consensus motif, 5'GGA(A/T)3' [13-17]. Both FLI1 and ERG proteins were shown to inhibit apoptosis [18]. Depending on the cellular environment, these factors can act as transcriptional activators, repressors, or both [19]. Of all the ETS factors, the ERG and FLI1 are the most closely related in terms of protein structure and have been extensively studied in the context of vascular homeostasis and remodeling [20,21].

Though ERG and FLI1 have similar structures, they have different cellular regulatory roles. ERG is mainly expressed in endothelial cells, megakaryocyte erythroid progenitor cells, T and B early lineage-committed cells, and myocardium [22]. On the other hand, FLI1 expression is mostly found in endothelial cells, hematopoietic progenitors and their mature derivatives (megakaryocytes, monocytes, erythroblasts, and natural killer cells), neural cells, and dermal fibroblasts [23].

ERG and FLI-1 in PAH Pathogenesis

Several in-vivo studies have shed more light on the biological roles of the ERG and FLI1 transcription factors especially in embryogenic development of cardiac structures and hematopoietic stem cells [24,25]. Notably, Looney and collaborators showed that the simultaneous small-interfering-

RNA (siRNA) knockdown of ERG and FLI1 in human pulmonary artery endothelial cells (HPAEC) and human pulmonary microvasculature endothelial cells (HPMEC) was associated with up-regulation of pro-inflammatory genes and interferon (IFN) pathway-related genes [26].

Additionally, dual knockdown of these factors had a significantly higher pro-inflammatory effect than obtained from knockdown of ERG or FLI1 alone. The study collaborators further compared the levels of endothelial ERG in the pulmonary vasculature of lung sections from patients with idiopathic PAH and scleroderma associated PAH to that of non-diseased lungs (healthy controls). The endothelium in normal lungs was found to have high levels of nuclear ERG compared to significantly lower levels to near absence of ERG in the endothelium of patients with diseased lungs. This suggests that down-regulation or loss of ERG in the HPAEC and HPMEC of humans is associated with systemic and pulmonary vasculopathy seen in PAH.

ETS Related Gene (ERG) and TGF- β 1/Smad3 pathway in the Pathogenesis of IPF

Smad3 has been shown to be a critical mediator in the pathogenesis of tissue fibrosis. In animal models, Smad3 signaling through phosphorylation and cytosol-nucleus translocation upregulated TGF- β 1-induced EMT processes and subsequent lung fibrosis [27,28]. Targeted suppression of the TGF- β 1/Smad3 pathway improved pulmonary fibrosis in rats¹⁰. Significantly elevated levels of TGF- β 1 has also been demonstrated in patients hitherto diagnosed with idiopathic pulmonary fibrosis [29].

In another study by Webb *et al.* [8], upregulation of ETS-related gene (ERG) promoted liver homeostasis by regulating canonical TGF β 1-SMAD signaling and promoting the SMAD1 pathway while repressing SMAD3 activity. Molecular ablation of ERG expression in this study further resulted in a SMAD3-dependent endothelial-to-mesenchymal transition (EMT) and spontaneous liver fibrogenesis. Decreased ERG expression was also shown to correlate with the degree of EndMT end-stage liver fibrosis. This study highlights ERG mediated fibrogenesis through the TGF β 1-SMAD pathway and the potential of ERG as a reliable biomarker in assessing EMT in tissue fibrosis.

Targeted Tumor necrosis factor-alpha (TNF- α) inhibition in PAH and IPF Treatment

Tumor necrosis factor-alpha (TNF- α) is a cell signaling protein (cytokine) produced by lymphocytes and macrophages. TNF- α mediates the immune response by attracting white blood cells to sites of injury and regulating inflammatory processes [30]. Dysregulation of TNF production has been implicated in a variety of autoimmune diseases. PAH and IPF patients demonstrate a high circulating levels of inflammatory cytokines especially TNF α , which also correlate with poor prognosis and survival [31,32].

Etanercept is a fusion protein produced by recombinant DNA that fuses the TNF receptor to the constant end of the immunoglobulin

(Ig) G1 antibody thereby reducing the inflammatory effects of TNF- α [33]. It is the most studied TN- α inhibitor in PAH treatment research. Studies have demonstrated that Etanercept administration prevented and reversed Monocrotaline (MCT) - induced PAH in experimental rats through the suppression of TNF- α expression and NF- κ B pathway [34-36]. Similar effects were also noted when Piguat and Vesin prevented Bleomycin and Silica induced lung fibrosis with the infusion of a TNF - alpha inhibitor in mice [37].

Etanercept as a Potential Treatment of PAH and IPF through ETS Expression and ERG mediation of TGF B1 / Smad3 Pathway

TNF- α inhibition may not be the only mechanism through which etanercept could be beneficial in PAH treatment. As already described, reduced expression of ERG and FLI1 transcription factors is associated with dysregulation of vascular homeostasis in HPAEC and HMPEC [26]. Likewise, upregulation of the TGF- β 1/Smad3 activate the cascade of lung fibroblasts proliferation and formation of aberrant epithelial cells leading to pulmonary fibrosis [38].

Notably, Neil *et al.* induced liver injury and fibrosis through administration of tetrachloric acid (CCL4) which caused a significant hyperplasia in smooth muscle actin (SMA), reduction of ERG expression and enhanced SMAD 3 phosphorylation [8]. The co-administration of etanercept prevented these changes and normalized endothelial ERG expression. Interestingly, etanercept was not able to prevent or correct the CCL4- induced injury in *Erg^{cEC-het}* mice which suggests that it potentially exerts its therapeutic effects through the endothelial ERG pathway in the prevention of fibrosis and restoration of vascular homeostasis.

Conclusion

Targeting the regulation of ERG, FLI1 and ERG-mediation of TGF- β 1/Smad3 signaling may be a promising therapeutic strategy in PAH and IPF. Etanercept appears to reduce endothelial inflammation and tissue fibrosis through increased expression of ERG and FLI1, promotion of ERG-mediated reduction of smad3 phosphorylation and suppression of TGF β 1/Smad3 signaling pathway. Tyrosine kinase (Ponatinib) has been shown to improve pulmonary fibrosis by inhibiting the TGF- β 1/Smad3 pathway invitro¹⁰. It is also tempting to speculate that FDA approved Smad3 inhibitors can be used as therapeutic agents through modulation of the functional roles of ERG and Fli-1 in PAH and IPF. ERG and Fli-1 protein levels may serve as early biomarkers for monitoring the health of blood vessels connected to the lung tissue in PAH and IPF patients. Similarly, serum levels of ERG and Fli-1 can be useful in monitoring treatment of patients with PAH and IPF as restoration to normal levels tend to correlate with positive response.

Conflict of Interest

We acknowledge that we have no conflict of interest or any financial interest that could influence our work.

Acknowledgements

We thank all the members of Reddy and Rao laboratories. This study was funded in part by the U.S. Army Medical Research and Materiel Command under W81XWH-08-1-0628, W81XWH-09-1-0236, W81XWH-10-1-0418 (Reddy, ESP) and the Georgia Cancer Coalition Distinguished Cancer Scholar award (Reddy, ESP and Rao, VN), U54/56 Morehouse School of Medicine/University of Alabama at Birmingham/Tuskegee University Partnership Grant (NIH 2U54CA118948, 3U54CA118638-05S1 to Dr Reddy), RCM1, U54 RR026137 and U54 MD007588.

References

1. Reingardiene D. Pulmonary arterial hypertension. *Medicina (Kaunas)*. 2007; 43(12):978-994.
2. Rosenkranz S. Pulmonary hypertension 2015: current definitions, terminology, and novel treatment options. *Clin Res Cardiol*. 2015. doi: 10.1007/s00392-014-0765-4.
3. Thompson K, Rabinovitch M. Exogenous leukocyte and endogenous elastases can mediate mitogenic activity in pulmonary artery smooth muscle cells by release of extracellular matrix-bound basic fibroblast growth factor. *J Cell Physiol*. 1996; 166(3):495-505. doi: 10.1002/(SICI)1097-4652(199603)166:3<495::AID-JCP4>3.0.CO;2-K.
4. Concepts C. Pathogenesis and pathophysiology. *Crit Care*. 1997:111-117. doi: 10.1056/NEJM199701093360207.
5. Raghu G, Rochwerf B, Zhang Y, Garcia CA, Azuma A, Behr J, et al. An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline: Treatment of Idiopathic Pulmonary Fibrosis. An Update of the 2011 Clinical Practice Guideline. *Am J Respir Crit Care Med*. 2015; 192(2):e3-e19. doi:10.1164/rccm.201506-1063st.
6. Wolters PJ, Collard HR, Jones KD. Pathogenesis of Idiopathic Pulmonary Fibrosis. *Annu Rev Pathol Mech Dis*. 2013; 9(1):157-179. doi: 10.1146/annurev-pathol-012513-104706.
7. Du Bois RM. Strategies for treating idiopathic pulmonary fibrosis. *Nat Rev Drug Discov*. 2010; 9(2):129-140. doi: 10.1038/nrd2958.
8. Neil P, Dufton, Claire R, Peghaire, Lourdes Osuna-Almagro, Claudio Raimondi, Viktoria Kalna, Abhishek Chuahan, et al. Dynamic regulation of canonical TGF β signalling by endothelial transcription factor ERG protects from liver fibrogenesis. *Nat Commun*. 2017. doi: 10.1038/s41467-017-01169-0.
9. Chinello P, Petrosillo N. Pharmacological treatment of HIV-associated pulmonary hypertension. *Expert Rev Clin Pharmacol*. 2016. doi: 10.1586/17512433.2016.1151785.
10. Qu Y, Zhang L, Kang Z, Jiang W, Lv C. Ponatinib ameliorates pulmonary fibrosis by suppressing TGF- β 1/Smad3 pathway. *Pulm Pharmacol Ther*. 2015; 34:1-7. doi: 10.1016/j.pupt.2015.07.004.
11. Reddy ES, Rao VN, Papas TS. The erg gene: a human gene related to the ets oncogene. *Proc Natl Acad Sci*. 2006; 84(17):6131-6135. doi: 10.1073/pnas.84.17.6131.
12. Rao VN, Papas TS, Reddy ES. erg, a human ets-related gene on chromosome 21: alternative splicing, polyadenylation, and translation. *Science*. 1987; 237(4815):635-639. doi: 10.1126/science.3299708. PMID 3299708.
13. Rao VN, Modi WS, Drabkin HD, Patterson D, O'Brien SJ, Papas TS, Reddy ES. The human erg gene maps to chromosome 21, band q22: relationship to the 8; 21 translocation of acute myelogenous leukemia. *Oncogene*. 1988; 3(5):497-500.
14. Prasad DDK, Rao VN, Shyam E, Reddy P. Structure and Expression of Human Fli-1 Gene. *Cancer Res*. 1992; 52(20):5833-5837.

15. Reddy ES, Rao VN D. *erg*, an ets-related gene, codes for sequence-specific transcriptional activator. *Oncogene*. 1991; 6(12): 2285–2289.
16. Siddique HR, Rao VN, Lee L, Reddy ES. Characterization of the DNA binding and transcriptional activation domains of the *erg* protein. *Oncogene*. 1993; 8(7): 1751–1755.
17. Rao VN, Ohno T, Prasad DD, Bhattacharya G, Reddy ES. Analysis of the DNA-binding and transcriptional activation functions of human Fli-1 protein. *Oncogene*. 1993; 8(8):2167-2173.
18. Yi H, Fujimura Y, Ouchida M, Prasad DD, Rao VN, Reddy ES. Inhibition of apoptosis by normal and aberrant Fli-1 and *erg* proteins involved in human solid tumors and leukemias. *Oncogene*. 1997; 14(11):1259–1268
19. Oikawa T, Yamada T. Molecular biology of the Ets family of transcription factors. *Gene*. 2003; 303(1-2):11-34. doi: 10.1016/S0378-1119(02)01156-3.
20. Kruse EA, Loughran SJ, Baldwin TM, Josefsson EC, Ellis S, Watson DK, et al. Dual requirement for the ETS transcription factors Fli-1 and Erg in hematopoietic stem cells and the megakaryocyte lineage. *Proc Natl Acad Sci*. 2009; 106(33):13814-13819. doi: 10.1073/pnas.0906556106.
21. Sato Y. Role of ETS Family Transcription Factors in Vascular Development and Angiogenesis. *Cell Struct Funct*. 2002; 26(1):19-24. doi: 10.1247/csf.26.19.
22. Birdsey GM, Shah AV, Dufton N, Reynolds LE, Osuna Almagro L, Yang Y, et al. The endothelial transcription factor *erg* promotes vascular stability and growth through Wnt/ β -catenin signaling. *Dev Cell*. 2015; 32(1):82-96. doi: 10.1016/j.devcel.2014.11.016.
23. Puzstaszeri MP, Seelentag W, Bosman FT. Immunohistochemical expression of endothelial markers CD31, CD34, von Willebrand factor, and Fli-1 in normal human tissues. *J Histochem Cytochem*. 2006; 54(4):385-395. doi: 10.1369/jhc.4A6514.2005.
24. Vijayaraj P, Le Bras A, Mitchell N, Kondo M, Juliao S, Wasserman M, et al. Erg is a crucial regulator of endocardial-mesenchymal transformation during cardiac valve morphogenesis. *Development*. 2012; 139(21):3973-3985. doi: 10.1242/dev.081596.
25. Spyropoulos DD, Pharr PN, Lavenburg KR, Jackers P, Papas TS, Ogawa M, et al. Hemorrhage, Impaired Hematopoiesis, and Lethality in Mouse Embryos Carrying a Targeted Disruption of the Fli1 Transcription Factor. *Mol Cell Biol*. 2002; 20(15):5643-5652. doi: 10.1128/mcb.20.15.5643-5652.2000.
26. Looney AP, Han R, Stawski L, Marden G, Iwamoto M, Trojanowska M. Synergistic role of endothelial *erg* and *fli1* in mediating pulmonary vascular homeostasis. *Am J Respir Cell Mol Biol*. 2017; 57(1):121-131. doi: 10.1165/rcmb.2016-0200OC.
27. Selman M, King J, Pardo A. Idiopathic pulmonary fibrosis: Prevailing and evolving hypotheses about its pathogenesis and implications for therapy. *Ann Intern Med*. 2001; 134(2):136-151. doi: 10.7326/0003-4819-134-2-200101160-00015.
28. Selman M. Role of Epithelial Cells in Idiopathic Pulmonary Fibrosis: From Innocent Targets to Serial Killers. *Proc Am Thorac Soc*. 2006; 3(4):364-372. doi: 10.1513/pats.200601-003tk.
29. Willis BC, Borok Z. TGF- β -induced EMT: mechanisms and implications for fibrotic lung disease. *Am J Physiol Cell Mol Physiol*. 2007; 293(3):L525-L534. doi: 10.1152/ajplung.00163.2007.
30. Barrett K, Chantry D, Turner M, Feldmann M, Gray PW. Cloning of human tumor necrosis factor (TNF) receptor cDNA and expression of recombinant soluble TNF-binding protein. *Proc Natl Acad Sci*. 2006; 87(19):7380-7384. doi:10.1073/pnas.87.19.7380.
31. Soon E, Holmes AM, Treacy CM, Doughty NJ, Southgate L, Machado RD, et al. Elevated Levels of Inflammatory Cytokines Predict Survival in Idiopathic and Familial Pulmonary Arterial Hypertension. *Circulation*. 2010; 122(9):920-927. doi: 10.1161/circulationaha.109.933762.
32. Oikonomou N, Harokopos V, Zalevsky J, Valavanis C, Kotanidou A, Szymkowski DE, et al. Soluble TNF Mediates the Transition from Pulmonary Inflammation to Fibrosis. *PLoS One*. 2006; 1(1):e108. doi: 10.1371/journal.pone.0000108.
33. Feldmann M, Maini RN. Erratum: TNF defined as a therapeutic target for rheumatoid arthritis and other autoimmune diseases (*Nature Medicine* (2003) 9 (1245-1250)). *Nat Med*. 2003; 9(11):1433. doi: 10.1038/nm1103-1433b.
34. Zhang LL, Lu J, Li MT, Wang Q, Zeng XF. Preventive and remedial application of etanercept attenuate monocrotaline-induced pulmonary arterial hypertension. *Int J Rheum Dis*. 2016. doi:10.1111/1756-185X.12304.
35. Zhang M, Wang Y, Wang Q, Zuo X, Xie W, Wang H. Monocrotaline-induced pulmonary arterial hypertension is attenuated by TNF- α antagonists via the suppression of TNF- α expression and NF- κ B pathway in rats. *Vascul Pharmacol*. 2012; 58(1-2):71-77. doi: 10.1016/j.vph.2012.07.006.
36. Koonin EV. The Origin at 150: is a new evolutionary synthesis in sight? *Trends Genet*. 2009; 25(11):473-475. doi: 10.1016/j.tig.2009.09.007.
37. Piguet PF, Ribaux C, Karpuz V, Grau GE, Kapanci Y. Expression and localization of tumor necrosis factor- α and its mRNA in idiopathic pulmonary fibrosis. *Am J Pathol*. 1993; 143(3):651-655.
38. Walton, Kelly L, Katharine E. Johnson, Craig A. Harrison. Targeting TGF- β mediated SMAD signaling for the prevention of fibrosis. *Frontiers in pharmacology*. 2017; 8(461).