

Risk Factors for Neonatal Depression in a Regional Public Hospital, Santos, Brazil, 2013

Marcele Brito de Souza¹, Sarah Warkentin², Laís Amaral Mais³ and Tulio Konstantyner^{4*}¹MD. Graduate Student in Pediatrics. Hospital Guilherme Álvaro (HGA), Santos, SP, Brazil²MSc. PhD Student of Department of Pediatrics, Universidade Federal de São Paulo (UNIFESP), Sao Paulo, SP, Brazil³MSc. PhD Student of Department of Pediatrics, Universidade Federal de São Paulo (UNIFESP), Sao Paulo, SP, Brazil⁴MD, MSc, PhD. Professor of Department of Pediatrics, Universidade Federal de São Paulo (UNIFESP) & Universidade Santo Amaro (UNISA), Sao Paulo, SP, Brazil

***Corresponding author:** Tulio Konstantyner, Discipline of Nutrology, Department of Pediatrics, Federal University of São Paulo. Rua Loefgreen, 1647, CEP: 04040-032, São Paulo, SP, Brazil; Tel/Fax: 55 (11) 5084-4538; E mail: tkmed@uol.com.br

Article Type: Research, **Submission Date:** 23 March 2017, **Accepted Date:** 8 June 2017, **Published Date:** 13 July 2017.

Citation: Marcelle Brito de Souza, Sarah Warkentin, Laís Amaral Mais and Tulio Konstantyner (2017) Risk Factors for Neonatal Depression in a Regional Public Hospital, Santos, Brazil, 2013. J.Paedi.Care.Inol 1(2): 7-11. doi: <https://doi.org/10.24218/jpci.2017.08>.

Copyright: © 2017 Marcelle Brito de Souza, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Neonatal depression leads to severe neurological sequelae with important impacts on quality of life. Most risk factors are sensitive to primary care actions, and this should be considered when developing prevention strategies. The aim was to identify the incidence and risk factors associated with neonatal depression in newborns at a Brazilian regional public hospital. A cross-sectional study of 784 live births with gestational ages of 22 weeks or more and without any major congenital anomalies was performed. Neonatal depression was defined as an Apgar score lower than seven at five minutes. Chi-square tests ($p < 0.05$) were used for the analysis of associations, and a logistic regression model was run to control the confounding factors. The neonatal depression rate was 2.2%. Multivariate logistic regression revealed that the presence of foul-smelling amniotic fluid (OR=14.2, $p=0.029$), prematurity (gestational age less than 37 weeks) (OR=37.5, $p < 0.001$) and the time of rupture of larger membranes (OR=8.7, $p=0.005$) were independently associated with neonatal depression. Risk conditions for perinatal infection and prematurity increase the likelihood of neonatal depression. Adequate prenatal care and guaranteed access to delivery services may reduce the occurrence of these neonatal depression risks and, consequently, may reduce infant morbidity and mortality.

Keywords: Asphyxia Neonatorum, Infant, Newborn, Risk Factors, Logistic Models, Primary Health Care.

Introduction

The main goal of neonatal care performed in the delivery room is to reduce infant morbidity and mortality, which promotes the survival of newborns (NBs) in the best possible functional conditions [1]. Neonatal depression is an injury that can lead to

hypoxic-ischaemic syndrome, and it affects NBs irrespective of gestational age.

Each year, approximately 4 million NBs worldwide present asphyxia, with one million developing severe sequelae and one million who do not survive [2,3]. Neurological sequelae impart a lower quality of life and higher costs to maintain the individual within the family and society [4,5].

In 1953, anaesthesiologist Virginia Apgar proposed a method to rapidly and accurately evaluate the birth conditions of children in their first minutes of life. The Apgar score considers five clinical parameters and ranges from zero to sixteen [6]. Apgar values between seven and ten in the fifth minute are considered normal. Values less than seven are inversely associated with an increased risk of neonatal morbidity and mortality, even though they do not necessarily predict neurological sequelae or any unfavourable outcomes [7,8].

Prematurity, low birth weight and birth asphyxia have been identified as the main factors associated with health problems at birth around the world. Maternal and gestational characteristics, such as infections, multiple births, hypertensive disease, smoking, illicit drug use, strenuous labour, low maternal schooling, black race, and a previous history of preterm birth, have been associated with a higher risk of low Apgar score (below seven) [9].

These factors, in an isolated or an aggregated way, may lead to poor health outcomes for children, especially those in less favoured populations that depend on the public health system. However, such factors are potentially modifiable by the control and prevention actions that are performed in primary health care activities. This fact represents a possibility for reducing the incidence of neonatal depression and its consequences.

In light of this fact, knowing the gestational characteristics and birth conditions of children is fundamental to subsidizing effective interventions. The aim is to adopt clinical practices that have been adjusted for populations with greater risk to reduce the morbidity and mortality associated with this undesirable hypoxic-ischaemic insult [10].

Thus, the present study aims to identify the incidence and risk factors associated with neonatal depression in a Brazilian regional public hospital.

Methods

This cross-sectional study was carried out from April 1st to September 30th, 2013, in the maternity ward and neonatal intensive care unit (NICU) of a public hospital located in Santos, Brazil. This hospital belongs to the Brazilian Unified Health System (SUS)/State Department of Health and is considered a reference centre for high risk pregnancy, attending, free of charge, the population of Baixada Santista, which has an average of 1,500 births per year. The study sample consisted of all live births in this period, regardless of the type of delivery, with gestational ages greater than or equal to 22 weeks. The exclusion criterion was the presence of a major congenital anomaly, such as hydrocephalus, diaphragmatic hernia, or kidney agenesis.

Data collection was performed by using the information available in medical records from the care team. Neonatal depression was defined as an Apgar score below seven in the fifth minute of life [9,11,12]. We used neonatal depression as an indicator of neonatal asphyxia. Characteristics related to the mother, gestational period, delivery and the NBs' first minutes of life were studied as risk factors. Among them, we studied premature amniorrhexis, meconium-stained amniotic fluid, preeclampsia, previous abortion, oligoamnion, intrauterine growth restriction (IUGR) and the presence of foul-smelling amniotic fluid (FSAF). In addition, we investigated the rupture time of membranes, type of delivery and presentation of the foetus, use of anaesthesia, number of previous deliveries, alcoholism, smoking and maternal educational level.

Adolescent pregnant women were under 20 years of age [13] and adequate prenatal care was a minimum of six visits performed during pregnancy [14]. NBs with a birth weight less than 2.500g were defined as low birth weight and gestational age less than 37 weeks was defined as prematurity.

The completed questionnaires were evaluated in terms of their internal consistency prior to data entry for analyses. Data were entered twice, and subsequent validation was conducted in order to correct errors. Stata statistical software package Version 14.0 (College Station, TX: StataCorp LP) was used to analyse the data.

Data consistency analysis and univariate descriptive statistics were performed for the continuous and categorical variables. A Pearson's chi-square test was used to compare categorical variables.

To adjust for confounding variables, a multivariate logistic regression analysis was performed using a "stepwise forward" technique in terms of odds ratio (OR) estimates. All variables with $p \leq 0.20$ were eligible for the final model and the results were statistically significant when $p \leq 0.05$.

This research received ethical approval from the Hospital Guilherme Álvaro – Santos ethics committee.

Results

Eight hundred and eleven births were recorded in the study period. Of these, 17 patients (2.1%) were not included because they were stillborn and ten (1.2%) were excluded because they had a major congenital anomaly at birth. Thus, 784 NBs composed the final sample.

The mean maternal age was 26 years, and 86% of mothers reported having a stable relationship with their partners. The majority (93.2%) of mothers reported some type of paid work; however, a third (28.1%) of mothers reported a study time equal to or less than eight years.

The rate of neonatal depression was 2.2% (17/782). Table 1 shows the incidences of the NB birth characteristics and comparisons

Table 1: Incidences of newborn birth characteristics and comparisons between the depressed non-depressed groups, Santos, Brazil, 2013

Characteristics (categorical variables)	n	Depressed	Non-depressed	p-value*
		%	%	
Low birth weight	782	76.5 (13/17)	12.5 (96/765)	<0.001
Male	780	47.0 (8/17)	53.0 (405/763)	0.633
Prenatal care	727	53.3 (8/15)	83.7 (596/712)	0.005
Instrumental delivery ^a	783	35.3 (6/17)	53.7 (411/766)	0.133
Adolescent mother	782	17.6 (3/17)	18.6 (142/765)	0.924
Meconium fluid	747	14.3 (2/14)	15.1 (111/733)	0.995
PPV with supplemental O ₂	774	93.7 (15/16)	6.4 (49/758)	<0.001
OTI	774	73.3 (11/15)	1 (8/759)	<0.001
Cardiac massage	774	13.3 (2/15)	0.13 (1/759)	0.001

^acaesarean or forceps; PPV: Positive pressure ventilation; OTI: Orotracheal intubation

*Chi-square test or Fisher's exact test

between groups with and without neonatal depression.

Table 2 presents the incidences and OR with respective confidence intervals for the risk factors associated with depression at birth. The presence of foul-smelling amniotic fluid at birth (OR=48.9 - 95% CI 6.2-318.2), prematurity (OR=26.8 - 95% CI 8.0-114.4) and low birth weight (OR=22.6 - 95% CI: 7.2-71.0) presented the greatest risks for neonatal depression.

The final logistic regression model (Table 3) shows the unadjusted and adjusted ORs for various risk factors of neonatal depression.

Discussion

The aim of this research was to identify the incidence and risk factors associated with neonatal depression. The incidence of this condition in the studied population was 2.2%, and the risk factors that were independently associated with neonatal depression were prematurity, presence of foul-smelling amniotic fluid, and longer membrane rupture times.

In a nationally representative Brazilian sample, Souza (2003) [15]

described a general incidence of neonatal depression of 2.2%, using the same definition as the present study used); however, there were regional variations. Thus, the incidence found in the present study corroborates the previously estimated rate in Brazil. The incidence in Brazil differs from worldwide neonatal depression rates, which range between three and six per 1000 live births. However, these rates have been estimated based on different diagnosis criteria, which makes comparisons between countries difficult [2].

The present study defined newborns with neonatal depression as those with Apgar scores below seven in the five-minute test. This cut-off has the potential to present good accuracy when it is applied in areas with limited available resources for a precise diagnosis [9,11]. This choice may have favoured sensitivity instead of specificity in the neonatal asphyxia diagnosis, which potentially resulted in an overestimation of the studied outcome.

Among the evaluated risk factors, we observed that eight newborns presented statistically significant associations with

Table 2: Incidences and odds ratios with respective confidence intervals (95% CI) of risk factors for depression at birth in Hospital Guilherme Álvaro, Santos, Brazil, 2013

Risk factors		OR	(CI 95%)	p-value*
Low Birth weight ^a	Yes	22.6	7.2 – 71.0	<0.001
	No			
Prematurity ^b	Yes	26.8	8.0 – 114.4	<0.001
	No			
Time of rupture of larger membranes (hours)	> 18	10.3	2.9 – 33.0	<0.001
	≤ 18			
Presence of FSAF	Yes	48.9	6.2 – 318.2	<0.001
	No			
Number of prenatal visits	< 6	0.2	0.1 – 0.6	0.001
	≥ 6			
General anaesthesia	Yes	14.3	1.5 – 138.0	0.002
	No			
Premature amniorrhexis	Yes	4.5	1.2 – 16.7	0.012
	No			
IUGR	Yes	3.6	0.8 – 16.7	0.075
	No			
Instrumental delivery ^c	Yes	0.5	0.2 – 1.3	0.142
	No			
Oligoamnion	Yes	1.6	0.2 – 12.4	0.656
	No			
Pre-eclampsia	Yes	0.7	0.1 – 5.1	0.695
	No			
Adolescent mother	Yes	0.9	0.3 – 3.3	0.924
	No			
Meconium fluid	Yes	0.9	0.2 – 4.2	0.929
	No			

^aBirth weight less than 2.500g, ^bGestational age less than 37 weeks, ^ccaesarean or forceps

OR: Odds Ratio; CI: Confidence Interval; FSAF: Foul-Smelling Amniotic Fluid; IUGR: Intrauterine Growth Restriction

*Chi-square test

Table 3: Final logistic regression model with unadjusted and adjusted odds ratios and their respective confidence intervals for risk factors for depression at birth, Santos, Brazil, 2013

Risk factors		Birth depression			
		OR [‡] (CI 95%)	p-value	OR [†] (CI 95%)	p-value
Prematurity ^a	Yes	26.8 (8.0-114.4)	<0.001	37.5 (7.7-182.2)	<0.001
	No	1.00			
Time of rupture of membranes (hours)	> 18	10.3 (2.9-33.0)	<0.001	8.7 (1.9-39.2)	0.005
	≤ 18	1.00			
FSAF	Present	48.9 (6.2-318.2)	<0.001	14.2 (1.3-151.7)	0.029
	Not present	1.00			

^aGestational age less than 37 weeks

OR: Odds Ratio; CI: Confidence Interval; FSAF: Foul-Smelling Amniotic Fluid

[‡]Unadjusted Odds Ratios; [†]Adjusted Odds Ratios (p<0.001)

neonatal depression: prematurity, low birth weight, the presence of foul-smelling amniotic fluid, a time of membrane rupture greater than 18 hours, a lack of prenatal care, intrapartum general anaesthesia, the presence of premature amniorrhexis, and a maternal history of syphilis. However, prematurity, the presence of foul-smelling amniotic fluid and a longer rupture of ovarian membranes were the only risk factors that remained in the multivariate final model; thus, these factors were independently associated with neonatal depression.

One of the main risk factors described in the literature for neonatal depression is a gestational age less than 37 weeks. Sousa [15] (2003) and Campos [16] (2010) observed that the presence of neonatal depression was inversely proportional to the gestational age. Similar results were found in our study, where prematurity presented an independent and positive association with neonatal depression. Body immaturity and difficulty transitioning from the foetal circulation to the neonatal circulation are the main factors that contribute to this association.

Klein *et al* [17] described that, during birth, an evaluation of vital signs with the Apgar score can be the first indicator of neonatal infection, with a correlation between a low five-minute test score and a higher risk of infection in the newborn. Longer times for membrane rupture and premature amniorrhexis predispose the mother and newborn to infectious diseases; these characteristics consequently lead to lower Apgar five-minute test scores. Chorioamnionitis, which has foul-smelling amniotic fluid as one of its clinical manifestations [18], also represents a risk for neonatal depression since it increases the risk of precocious neonatal sepsis. In our study, we observed that membrane rupture times greater than 18 hours and the presence of foul-smelling amniotic fluid showed independent and statistically significant associations with neonatal depression.

Lower birth weight is described by Sarinho *et al* [19] as a determining factor of neonatal mortality. In Brazil, according to Guimarães and Melendez [20], the lower birth weight rate is 9.2%, with possible variations by region. In the present study,

the lower weight birth rate was 13.9%, and it was significantly associated with neonatal depression; similar results were also found by Cunha *et al* [12] in his research in a maternity ward in Rio de Janeiro in 2004.

In parallel, it was observed that most pregnant women had at least six prenatal consultations (82.9%), as has been observed in other reference hospitals [21]. This consultation rate can be explained by the fact that high risk pregnancies require higher care and result in higher adhesion to prenatal care regimens. In this study, having more prenatal consultations was shown as a protective variable for neonatal depression; this result corroborates what those described by Rosa *et al* [22] in 1999. However, this association was not significant in the multivariate model, since prematurity and low birth weight are highly associated with a lack of or fewer prenatal consultations [23-25].

The present study identified prematurity, the presence of foul-smelling amniotic fluid and membrane rupture times greater than 18 hours as statistically significant risk factors that were independently associated with neonatal depression. This finding was further validated by including explanatory factors that have the potential to influence neonatal depression in the multivariate analysis; our approach provided a broad perspective, which is necessary for events that are triggered by multiple risk factors [26].

However, it is important to mention that the present study did not include other potential risk factors for birth depression, such as cardiotocogram abnormalities and maternal fever. Therefore, the internal validity of the results must be considered with caution.

In addition, it is noteworthy to mention that it is very important to prevent perinatal depression by systematically examining prenatal risk factors and giving interventions to newborns with risk factors, especially those with multiple risk factors, as the risk of presenting depression increases when more risk factors are present [27].

In this context, the results reinforce the need for direct adoption of preventive measures in prenatal care, with improvements in

primary health care for pregnant Brazilian women. The evidence from these findings indicates that, during the elaboration and execution of control and prevention strategies for neonatal depression, health policies should consider the effectiveness of prenatal consultations, which promote better care for pregnant women, especially those who have higher risks of premature birth and infectious diseases during pregnancy.

Acknowledgement

We would like to thank the participating mothers and the team of the Maternal Child Specialised Reference Hospital in Santos (Hospital Guilherme Álvaro), where the study was conducted.

References

1. Tragante CR, Ceccon MEJ, Falcão MC. Desenvolvimento dos cuidados neonatais ao longo do tempo. *J Pediatr (Rio J)*. 2010; 32(2):121-130.
2. Cruz ACS, Ceccon MEJ. Prevalência de asfixia perinatal e encefalopatia hipóxico-isquêmica em recém-nascidos de termo considerando dois critérios diagnósticos. *Rev Bras Cresc Desenvol Hum*. 2010; 20:302-316.
3. Majeed R, Memon Y, Majeed F, Shaikh NP, Rajar UDM. Risk factors of birth asphyxia. *J Ajub Med Coll Abbottabad*. 2007; 19(3):67-71.
4. Macdonald HM, Mulligan JC, Allen AC, Taylor PM. Neonatal Asphyxia I. Relationship of obstetric and neonatal complications to neonatal mortality in 38,405 consecutive deliveries. *J Pediatr*. 1980; 96(5):898-902.
5. Rosa IRM, Marba STM. Asfixia neonatal: incidência e mortalidade neonatal em 5424 partos consecutivos. Relação com o peso e idade gestacional. *Anais do XIV Congresso Brasileiro de Perinatologia e XI Reunião de Enfermagem Perinatal*, 1994, São Paulo.
6. Apgar V. The newborn (Apgar) scoring system. Reflections and advice. *Pediatr Clin North Am*. 1966; 13(3):645-650.
7. Casey BM, McIntire DD, Leveno KJ. The continuing value of the Apgar score for the assessment of newborn infants. *N Engl J Med*. 2001; 344(7):467-471.
8. American Academy of Pediatrics Committee on Fetus and Newborn: Use and abuse of the Apgar score. *Pediatrics*. 1986; 78(6):1148-1149.
9. Odd D, Doyle P, Gunnell D, Lewis G, Whitelaw A, Rasmussen F. Risk of low Apgar score and socioeconomic position: a study of Swedish male birth. *Acta Paediatrica*. 2008; 97(9):1275-1280. doi: 10.1111/j.1651-2227.2008.00862.x.
10. Horbar JD, Plsek PE, Schriefer JA, Leahy K. Evidence-based quality improvement in neonatal and perinatal medicine: the neonatal intensive care quality improvement collaborative experience. *Pediatrics*. 2006; 118:557-64.
11. Oswyn G, Vince JD, Friesen H. Perinatal asphyxia at Port Moresby General Hospital: a study of incidence, risk factors and outcome. *PNG Med J*. 2000; 43(1-2):110-120.
12. Cunha AA, Fernandes DS, Melo PF, Guedes MH. Fatores Associados à asfixia perinatal. *Rev Bras Ginecol Obstet*. 2004; 26(10):799-805.
13. World Health Organization, WHO. Child and adolescent health and development: progress report 2006-2007 [text on the internet]. Geneva; 2008. Available at: <http://www.who.int/maternal_child_adolescent/documents/9789241596497/en/>.
14. Brasil. Ministério da Saúde. Datasus. Informações de saúde. Estatísticas vitais: mortalidade e nascidos vivos. Available at: <<http://tabnet.datasus.gov.br/cgi/deftohtm.exe?sinasc/cnv/nvuf.def>>.
15. Souza FMS. Fatores associados à asfixia perinatal no Brasil: estudo populacional com base no Sistema de Informações de Nascidos Vivos. PhD thesis. Instituto Fernandes Figueira, Fundação Oswaldo Cruz. Rio de Janeiro, RJ, 2003.
16. Campos NG. Prevalência de asfixia perinatal e fatores associados em Fortaleza - Ceará. Masters dissertation. Centro de Ciências da Saúde, Universidade Estadual do Ceará. Fortaleza, CE, 2010.
17. Klein JO, Marcy SM. Bacterial sepsis and meningitis. In: Remington JS, Klein JO. Infectious diseases of the fetus and newborn infant. 3th ed. Philadelphia: W.B. Saunders; 1990. p. 610-644.
18. Araújo MCK, Feferbaum R, Vaz FAC, Ramos JLA. Infecção neonatal, rotura prematura de membranas amnióticas e corioamnionite. *Pediatrics (São Paulo)*. 1994; 16(3):95-101.
19. Sarinho SW, Filho DAM, Silva GAP, Lima MC. Fatores de risco para óbitos neonatais no Recife: um estudo caso-controle. *J Pediatría (Rio J)*. 2001; 77:294-298.
20. Guimarães EA, Velasquez-Melendez G. Low birth weight determinants from the Born Alive National Surveillance System in Itáúna, Minas Gerais. *Rev Bras Saude Mater Infant*. 2002; 2(3):283-290.
21. Santos GHN, Martins MG, Sousa MS, Batalha SJC. Impacto da idade materna sobre os resultados perinatais e via de parto. *Rev Bras Ginecol Obstet*. 2009; 31(7):326-334.
22. Rosa IRM, Marba STM. Fatores de risco para asfixia neonatal em recém-nascidos com peso acima de 1000 gramas. *J Pediatr (Rio J)*. 1999; 75(1):50-54.
23. Sabiri N, Kabiri M, Razine R, Barkat A. Risk factors leading to pre-term births in Morocco: a prospective study at the maternity Souissi in Rabat. *Pan Afr Med J*. 2015; 10; 22:21. doi: 10.11604/pamj.2015.22.21.5100.
24. Xaverius P, Alman C, Holtz L, Yarber L. Risk Factors Associated with Very Low Birth Weight in a Large Urban Area, Stratified by Adequacy of Prenatal Care. *Matern Child Health J*. 2016; 20(3):623-629. doi: 10.1007/s10995-015-1861-4.
25. Rani S, Chawla D, Huria A, Jain S. Risk factors for perinatal mortality due to asphyxia among emergency obstetric referrals in a tertiary hospital. *Indian Pediatr*. 2012; 49(3):191-194.
26. Kleinbaum DG, Kupper LL, Nizam A, Muller KE. Applied regression analysis and other multivariable methods. 4th ed. Duxbury: Brooks/Cole; 2008.
27. Chen ZL, He RZ, Peng Q, Guo KY, Zhang YQ, Yuan HH, et al. Prenatal risk factors for neonatal asphyxia: how risk for each? *Zhongguo Dang Dai Er Ke Za Zhi*. 2009; 11(3):161-165.